

# SYMPTOMS OF RADIO FREQUENCY ATTACK!

Do you see your self here on this list?

<http://www.md-spec.com>

## 1st Stage Signs

Both children and adults need to carefully attempt to sense as they are drifting off to sleep to determine whether they are feeling a shock as one drifts off to sleep, or seeing lights as one drifts off to sleep. Due to the difficulty of remembering such occurrences as one drifts off to sleep

or wakes up. It is recommended that a piece of paper and pencil be held in the hands on the chest where the person could make a mark when such instances occur.

Accompanying activity of sabotage of business and or personal life, strange people in area, strange reactions from people, items moved or missing in home.

Unaccountable increased heart rate.

Nervousness.

Irritability.

Aggravation reactions.

Erection dysfunction.

Compressing the vertebrae noted upon waking.

Sensitivity to sound.

Depression.

Minor spasms and Charlie horses.

Inability to concentrate.

Loss of memory.

Night sweats.

Sensitive ear to touch.

Upon waking finding that the muscles of the back are tingling as if electrified.

Ringling in one or both ears and or hear tone bursts (Tinnitus)

GERD (Difficulty in Swallowing)

Arthritis.

Eye Sight becomes blurred.

Teeth snap together when drowsy.

Loss of sleep.

Nails become wavy.

Water weight and cellulite accumulations on the upper thigh

and buttocks from cellular fluids displaced generally

## 2st Stage Signs

Burning sensation on the skin.

Loss of hair.

Feeling of temporary heating of the head. (demodulating RF effect)

Rashes.

Narcoleptic reactions, sleepiness.

Sleep only after exhaustion.

Virtual insomnia.

Miscarriage.

Gaunt face.

Facial wrinkles.

Losing skin turgor.

Hair breaks from rapid microwave heat on hair causing increase fracturing of hair shafts from expansion of water via humidity.

Loosing Control (unexplained anger)

Heating of body. (Body feeling as if it is being cooked)

Waking after sleep feeling as though you did not sleep, children's

symptoms can manifest themselves as ADD and Hyperactivity

## 3rd Stage effects from exposure

Loss of coordination.

Accidents from sleeplessness.

Damage to eyesight.

Atrophy of the muscles.

Heart valve damage.

Loss of weight and/or weight gain.

Nausea.

Sensitivity to sound.

Decreased dexterity.

Seizure.

Choking.

Vivid dreams.

Lost time. (Alzheimer type symptoms)

Change of mood.

Heart Attack.

The personality becomes quiet, a lack of thought takes place.

from the head and upper body especially in women, where those fluids find ready redistribution in those areas. Changes in the tone of the internal voice. Waking up in extreme pain, and arthritis condition. Spontaneous tearing without emotional thinking. Mild pressure in the head, a cloudy feeling. Increased need to urinate during the night. Feeling puffs of air on the face and back, such may be a demodulation of energy on the skins surface. Sensation that blood is trickling into localized areas of the brain and other parts of the body, this feeling is akin to the impression that these areas were devoid of blood. Unaccountable increased heart rate just before drifting off to sleep causing the person to wake up.

Decrease in mental activity.  
Limbs jerk typically as one is trying to sleep.  
Weakness.  
Lethargy or hyperactivity.  
Forgetfulness.  
Constipation.

4th Stage  
Syndromes.  
Disease.  
Insanity.  
Heart Attack.  
Stroke.  
Death.

<http://www.md-spec.com>